


Microbiology of Human Spacecraft Environments



C. Mark Ott, PhD

Microbiology Laboratory

NASA Johnson Space Center, Houston, TX







Microbiology and Space Missions

- Astronaut health
- Spaceflight foods
- Vehicle habitat
 - Life support systems
 - Trash and human waste containment
 - Astronaut hygiene areas
- Planetary protection
 - How do we protect the crew and extraterrestrial environments in our search for life on other planets?



These priority areas for microbiology drive vehicle design, microbial monitoring, and operational research efforts



Astronaut Health

- Diagnosis is often based on symptomology (*e.g.*, headache, rash, dry hacking cough, diarrhea)
- Examples of infectious diseases during spaceflight missions include upper respiratory infections, urinary tract infections, ear infections, herpes zoster, rashes and skin disorders, and gastroenteritis.
- Survey of Space Shuttle missions (STS-1 through STS-89) indicated infectious disease accounted for 1.4% of all medical events (not including skin and subcutaneous tissue)
- Evidence of immune system dysfunction

Why do exceptionally healthy astronauts in a pristine environment still contract infectious diseases?





Spaceflight Food


- Several categories
 - Fresh foods
 - Irradiated foods
 - Thermostabilized foods
 - Non-thermostabilized foods
- Non-thermostabilized foods
 - Includes dehydrated foods, tea, food bars, and cheese
 - Samples from each production lot are tested
 - Lots have been disqualified after detection of microbes, including the bacteria *Salmonella enterica* Typhimurium, *Staphylococcus aureus*, *Klebsiella pneumoniae*, and fungi *Aspergillus fumigatus*, and *Aspergillus flavus*.





The Future of Spaceflight Food

- The VEGGIE spaceflight experiment grew lettuce on the International Space Station (ISS)
 - Can the crew eat it?
- Probiotics are being evaluated as immune boosters
 - What organisms do we monitor for safety?
- We are planning to prepare and cook more food during flight
 - How do we control contamination during processing?





Vehicle Habitat - Spacecraft Design



- Vehicle surfaces
 - Use of materials that are not conducive to microbial growth
- Vehicle air
 - Use of High Efficiency Particulate Air (HEPA) filters
- Potable water recovery system
 - Potable water can be brought on a mission, or recycling humidity condensate and/or urine may be more efficient
 - On ISS, recycled water experiences a catalytic oxidizer (267°F for 10 minutes); iodine disinfection; and passes through a 0.2-micron filtration



Vehicle Habitat - Operations

- Health Stabilization Program
 - Preflight quarantine to decrease the risk of infectious disease
 - Employs other prevention techniques, such as raising infectious disease awareness
- Preflight disinfection and microbial monitoring
- Inflight cleaning and remediation
 - Regular housekeeping
 - Disinfectant wipes for contaminated surfaces
 - High concentration iodine solution to disinfect potable water systems





Contamination Potential



Spacecraft are complex



Trash and waste control



Astronaut activities, such as eating and hygiene




Microbial Monitoring on the ISS


Surfaces



Air





Water



Quantified in-flight and returned to JSC for identification




Microbial Monitoring of Water





Advances in Microbial Monitoring

- DNA sequencing in space
 - First performed in 2016 using Oxford Nanopore's MinION technology
- In-flight identification of environmental samples
 - Three microbial colonies from a media plate used for ISS surface sampling were processed and sequenced on ISS.
 - The MinION accurately identified the three isolates that were selected.
- Following up with Swab to Sequencer direct analysis



Trade names and trademarks are used in this report for identification only. Their usage does not constitute an official endorsement, either expressed or implied, by the National Aeronautics and Space Administration.



Environmental Health

Microbial monitoring by NASA and others indicate ISS environmental flora reflect human-associated microorganisms commonly found in terrestrial homes.



- Media-based monitoring occasionally identifies opportunistic bacterial pathogens, such as *Staphylococcus aureus* and *Bacillus cereus*. No methicillin resistant *S. aureus* (MRSA) has been identified.
- *Enterobacter* and *Enterococcus* species are occasionally identified throughout the ISS.



Environmental Health

- The most prevalent fungal genera are *Aspergillus* and *Penicillium*. *A. flavus*, *A. niger*, and *A. fumigatus* have been identified from spaceflight samples. *Stachybotrys chartarum* has been isolated preflight.
- Samples from potable water (for drinking and hygiene) indicate common water-borne bacterial species.




As ISS has many visiting vehicles with astronauts, “new” bacterial and fungal isolates are regularly identified.




Environmental Anomalies

- Russian Mir Space Station (1986-2001)
 - Samples of free-floating condensate caused by power failures and problems with temperature control revealed high microbial diversity, including *Escherichia coli*, *Serratia marcescens*, *Legionella* species, and protozoa
- International Space Station (1998-present)
 - Surface contamination is often associated with uncontrolled water
 - Air contamination and unpleasant odors do occur; however, the sources are often unknown. These events have been transient and/or handled by the HEPA filtration system.
 - Biofilm formation in ISS water lines have caused clogged lines and pump failure.





ISS Environmental Anomalies






Microbiological Research in Space

- Multiple spaceflight experiments over the past 50 to 60 years have demonstrated unexpected microbial responses when microorganisms are cultured during spaceflight, including:
 - Antibiotic resistance
 - Biofilm formation
 - Growth rates
 - Virulence
 - Gene expression
- Rotating Wall Vessel (RWV) bioreactor
 - Fluid movement in the reactor simulates several aspects of the microgravity environment
 - Provides predictive and follow-up studies for spaceflight experiments



The Fluid Processing Apparatus (FPA), an example of in-flight hardware




The low fluid shear culture conditions has prompted the term Low Shear Modeled Microgravity (LSMMG) environment




MICROBE: *Salmonella* Response to Spaceflight Culture

PI: Dr. Cheryl Nickerson, Arizona State University

Flight Sample



Ground Control



- In 2006, the MICROBE experiment compared the characteristics of *Salmonella* Typhimurium grown in space to otherwise identical cultures grown on the Earth.

- First to identify alterations in microbial virulence in a spaceflight grown microorganism
- First transcriptomic study of a microorganism grown in space
- First identification of a spaceflight molecular response mechanism
- First to identify novel changes in biofilm production



Astronaut Microbiome

PI: Dr. Hernan Lorenzi, J. Craig Venter Institute



- First comprehensive study of the microbiomes of nine astronauts on ISS
- Samples included skin sites, nostrils, fecal samples collected preflight, in-flight, and post-flight
- Composition of the intestinal microbiota became more similar across astronauts in space
- Alterations in the skin microbiome that might contribute to the high frequency of skin rashes
- Set the baseline for astronaut microbiome studies with low sample size



Selected Current Studies

PI: Dr. Mark Ott, NASA

Co-PI: Dr. Cheryl Nickerson, Arizona State University

- Determine if growth in the spaceflight analogue (RWV) environment increases virulence in other bacteria, five pathogenic species are being investigated to characterize changes in stress response, adhesion/invasion of 3-D tissue culture models, and animal testing if warranted.

PI: Dr. Cheryl Nickerson, Arizona State University

- Investigate the combined effect of spaceflight analogue (RWV) culture with low dose radiation to determine if the combination synergistically increases the potential infection risk.

PI: Dr. Bob McLean, Texas State University

Co-PI: Dr. Cheryl Nickerson, Arizona State University

- Investigate polymicrobial biofilm development by *Pseudomonas aeruginosa* and *Escherichia coli* during spaceflight to identify changes in biofilm architecture, disinfection, and corrosion potential during spaceflight.



QUESTIONS???